

Must Hypothalamic Neurosecretory Function Cease for Brain Death Determination? No

The UDDA Revision Series

Panayiotis N. Varelas, MD, PhD

Neurology® 2023;101:137-139. doi:10.1212/WNL.0000000000207336

Correspondence

Dr. Varelas
panayiotis.n.varelas@gmail.com

The Uniform Determination of Death Act of 1980,¹ which provides the statute of death in the United States, states,

1. [Determination of Death]. An individual who has sustained either (1) irreversible cessation of circulatory and respiratory functions or (2) irreversible cessation of all functions of the entire brain, including the brain stem, is dead. A determination of death must be made in accordance with accepted medical standards.

Most of the attention has been placed on the word “entire.” The reason is that in other countries, such as the United Kingdom, the entire brain criterion is not required and instead the brainstem criterion is deemed adequate.² In the United States, the word “entire” has also generated debate, with a small minority arguing that when used to the letter, up to half of brain dead (BD) patients are erroneously diagnosed as dead and, in fact, when heart-beating organ donation follows, these mistakenly diagnosed dead patients may be considered homicide victims.³ I will try to refute this proposition.

The rationale behind this serious accusation is that not all BD patients lose pituitary activity. The reason for any residual pituitary activity may be traced to anatomical variances that theoretically could allow some perfusion of these basal brain structures.^{4,5} For example, central diabetes insipidus (DI) was encountered in 65% of 266 BD patients included in a recent large single-center study.⁶ In review articles, 50%–60% of adult and 52% of pediatric BD patients were diagnosed with DI.^{3,7} The authors of these review articles speculate that hypothalamic and pituitary *functions* were still present. This argument is problematic for many reasons: First, not every study used rigorous diagnostic criteria for central DI. Polyuria is an easily noticed feature, but not pathognomonic in patients treated with diuretics, such as mannitol. Rising serum sodium is also not indicative of DI when these patients receive osmotic agents that have an uplifting effect on sodium. If hypotension occurs and vasopressin infusion is used, this external hormone will not allow DI to clinically manifest and may be confused with inherent hormone production. Similarly, if the peripheral organ (the kidney) has decreased function because of an acute or chronic process, clinical signs of DI will not be observed. Although this association has not been rigorously assessed, in a small sample of pediatric patients,⁸ in the aforementioned study,⁶ those patients who did not develop DI had statistically significant higher blood urea nitrogen and creatinine levels (implying kidney dysfunction) than those who did (*Varelas, unpublished data*). Therefore, it is possible that a higher percentage of posterior pituitary or hypothalamic loss of function could be present because of the “masking” effect on DI.

Similar to vasopressin, other hypothalamic hormones may or may not be reduced in BD patients, but because they do not lead to an obvious clinical sign (such as polyuria), they are not regularly measured nor reported.³ Although there is a paucity of autopsy data on pathologic injury of the hypothalamus/pituitary gland in BD patients, the few studies that exist are revealing. In the National Institute of Neurological and Communicative Disorders and Stroke Collaborative “nonintervention” study, 226 of 459 patients who died had brain autopsies. Although the thalamus/hypothalamus was one of the areas examined, the pituitary gland was not. The diencephalon had damage in 85% of the cases, but the hypothalamic and

RELATED ARTICLE

Contemporary Issues in Practice, Education, & Research

Must Hypothalamic Neurosecretory Function Cease for Brain Death Determination? Yes: The UDDA Revision Series

Page 134

From the Albany Medical College, NY.

Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

Glossary

ADH = antidiuretic hormone; BD = brain dead; DI = diabetes insipidus.

periventricular regions contained normal cells among necrotic areas.⁹⁻¹¹ In another study of 60 BD patients, the central part of the anterior pituitary lobe showed autolysis, but the marginal part was spared if the autopsy was performed in BD patients of less than 10 days.¹² Similarly, 28 patients with structural injuries were diagnosed as BD by clinical criteria and with negative intracranial blood flow and were maintained on ventilatory support until cardiac arrest ensued.⁵ Duration of the post-BD diagnosis period played a role in the pathologic findings of the 12 autopsied patients. Extensive necrosis of the brain including the hypothalamus was observed only in patients maintained for longer than 3 days. The hypothalamus was completely necrotic in cases with brain death duration of >17 hours. More revealing was the fact that adrenocorticotropic hormone and thyroid stimulating hormone–positive substances were clearly recognized in the necrotic cells, implying that these hormones may be maintained even in dead cells. The posterior pituitary gland showed little or no change in 11 of 12 autopsies. Of great importance, however, was the observation that intrinsic antidiuretic hormone (ADH) secretion decreased sharply with the occurrence of BD, implying cessation of supraoptic and paraventricular nuclei function, but plasma levels were detectable and ADH-positive cellular granules were observed long time after BD, indicating “leak” of ADH to the periphery and thus masking of clinical DI. This would *mimic osmoregulation* (which, in reality, is not present), as even the detractors of entire brain function loss, eventually admit.⁷ Finally, for other hypothalamic hormones, such as gonadotropin-releasing hormone and luteinizing hormone-releasing hormone, that are detected in the serum long after the hypothalamus becomes completely necrotic and without having any correlation with levels of luteinizing hormone, follicle-stimulating hormone, or human growth hormone, an extracerebral source is the likely explanation.⁵

The study by Sugimoto et al. is remarkable for another reason. It introduces the time concept for these pathologic changes and shows that as time passes, even initially preserved areas in the pituitary gland lose cellular integrity and become necrotized. More recent pathologic studies may not have found these alterations, including the “respirator brain,” because of the rapid advance to organ harvesting or withdrawal from support, which has shortened the time from BD to autopsy. For example, in a more recent study of 41 BD patients, all had cardiac arrest after BD declaration within 36 hours, with 29% in <12 hours. The pituitary gland showed moderate-to-severe neuronal loss in 45% of the 16 pituitary glands examined.¹³

One may then ask the following question: Even in rare patients with hypothalamic neurosecretory activity, how relevant is this? Are these BD patients not dead, that is, patients

who can “survive?” In the discussion regarding the Construction of the Statute (pages 75–79), the President’s Commission focused on the definition of “individual,” “functions,” and “dead,” leaving out any meaningful discussion on the keyword “irreversible.”¹⁴ Moreover, the Commission makes an important distinction between brain functions and cellular activities, clarifying that electrical or metabolic *activities* at an individual cell level (e.g., hormones) or even groups of cells, *can still be present* for a period of time, but unless organized and directed, they are considered “irrelevant” in judging whether the organism, as opposed to its components, is “dead.”

Irreversible cessation of a vital function means that once it ceases, it cannot be restored with available technology because doing so is impossible.¹⁵ Another way of saying this is that irreversibility is loss of brain function that cannot resume spontaneously and cannot be resumed through technological intervention. Key here is the understanding that although the function can be replaced by technology (e.g., blood circulation by artificial heart, breathing by ventilator), the *organ function cannot be restored by the organism* (no circulation without a heart, no breathing without an alive medulla oblongata). Consciousness or brainstem reflexes cannot be restored spontaneously or with interventions in BD patients with severe, diffuse cellular-level injury and no intracranial circulation. However, most important is that these patients *cannot spontaneously restore cardiorespiratory function*. This loss of the lower medullary function, as clinically noticed by the cardiocirculatory collapse and the apnea testing in patients who have additional diffuse injury to the rest of the brain, are in deep coma and with loss of all the other brainstem reflexes, cannot be restored either by the patient or by our current technology (no artificial brain or transplanted brain are available), *and is irreversible*. And this despite allowing years to pass, as in those rare BD patients chronically maintained on ventilators; none regained spontaneous respiratory function.¹⁶ This is a completely different situation than traumatic upper cervical spine injury with inability to breathe, where consciousness, brainstem reflexes, and supratentorial connectivity are preserved. In the former, no hypothalamic/pituitary action can counteract the profound loss of integrative function of the rest of the brain. When these BD patients (who are not organ donors) are disconnected from technological support, their blood pressure plummets, the pulse disappears, and the heart stops beating within minutes. These patients reach the same end point and look exactly alike to those who lose cardiorespiratory function after cardiac arrest. No breathing means no heart beat and no circulation, exactly like no heart beat means no circulation and no breathing, all these despite any potential hypothalamic/pituitary secretions. BD patients are dead the same way that everybody became dead since the dawn of man and before ventilators and intensive care units were invented.

In conclusion, residual hypothalamic neurosecretory function is highly unlikely (as compared with activity, which may be present in a small percentage of BD cases). Requirement for cessation of this activity does not add to the declaration of brain death and is irrelevant because these patients with irreversible cessation of all other functions of the brain cannot survive, similar to patients without cardiopulmonary function. These cardiopulmonary dead patients may also have hypothalamic-pituitary activity for a few minutes until their cells become irreversibly damaged, but nobody has ever confirmed such activity in a pulseless, nonbreathing dead patient, because this is irrelevant to their death.

Study Funding

No targeted funding reported.

Disclosure

The author reports no disclosures. Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures.

Publication History

Received by *Neurology* September 18, 2022. Accepted in final form March 7, 2023. Submitted and externally peer reviewed. The handling editor was Editor-in-Chief José Merino, MD, MPhil, FAAN.

Appendix Author

Name	Location	Contribution
Panayiotis Nicolaou Varelas, MD, PhD	Albany Medical College, NY	Drafting/revision of the manuscript for content, including medical writing for content; Study concept or design

References

1. National Conference of Commissioners on Uniform State Laws. *Uniform Determination of Death Act* [online]. Accessed December 20, 2022. [higherlogicdownload.s3-external-1.amazonaws.com/UNIFORMLAWS/4d19d096-be64-3c0f-ae71-a514b64c06a6_file.pdf?AWSAccessKeyId=AKIAVRD07IEREB57R7MT&Expires=1671548456&Signature=OBSiMl019H6l8jDnOsThH2MomPg%3D](https://www.amazonaws.com/UNIFORMLAWS/4d19d096-be64-3c0f-ae71-a514b64c06a6_file.pdf?AWSAccessKeyId=AKIAVRD07IEREB57R7MT&Expires=1671548456&Signature=OBSiMl019H6l8jDnOsThH2MomPg%3D).
2. Academy of Medical Royal Colleges. *A Code of Practice for the Diagnosis and Confirmation of Death* [online]. Accessed December 20, 2022. [aomrc.org.uk/reports-guidance/ukdec-reports-and-guidance/code-practice-diagnosis-confirmation-death/](https://www.aomrc.org.uk/reports-guidance/ukdec-reports-and-guidance/code-practice-diagnosis-confirmation-death/).
3. Nair-Collins M, Joffe AR. Frequent preservation of neurologic function in brain death and brainstem death entails false-positive misdiagnosis and cerebral perfusion. *AJOB Neurosci*. 2021;1-14.
4. Blumenfeld H. *Neuroanatomy Through Clinical Cases*, 2nd ed. Sinauer Associates; 2010.
5. Sugimoto T, Sakano T, Kinoshita Y, Masui M, Yoshioka T. Morphological and functional alterations of the hypothalamic-pituitary system in brain death with long-term bodily living. *Acta Neurochir*. 1992;115(1-2):31-36.
6. Varelas PN, Rehman M, Mehta C, et al. Comparison of 1 vs 2 brain death examinations on time to death pronouncement and organ donation: a 12-year single center experience. *Neurology*. 2021;96(10):e1453-e1461.
7. Nair-Collins M, Northrup J, Olcese J. Hypothalamic-pituitary function in brain death: a review. *J Intensive Care Med*. 2016;31(1):41-50.
8. Finfer S, Bohn D, Colpitts D, Cox P, Fleming F, Barker G. Intensive care management of paediatric organ donors and its effect on post-transplant organ function. *Intensive Care Med*. 1996;22(12):1424-1432.
9. An appraisal of the criteria of cerebral death. A summary statement. A collaborative study. *JAMA*. 1977;237(10):982-986.
10. National Institute of Neurological and Communicative Disorders and Stroke. *The NINCDS Collaborative Study of Brain Death*. U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health, National Institute of Neurological and Communicative Disorders and Stroke; 1981.
11. Walker AE, Diamond EL, Moseley J. The neuropathological findings in irreversible coma: a critique of the "respirator brain." *J Neuropathol Exp Neurol*. 1975;34(4):295-323.
12. Ujihira N, Hashizume Y, Takahashi A. A clinico-neuropathological study on brain death. *Nagoya J Med Sci*. 1993;56(1-4):89-99.
13. Wijdicks EF, Pfeifer EA. Neuropathology of brain death in the modern transplant era. *Neurology*. 2008;70(15):1234-1237.
14. President's Commission for the Study of Ethical Problems in Medicine and Biomedical and Behavioral Research. *Defining Death: A Report on the Medical, Legal and Ethical Issues in the Determination of Death*. U.S. Government Printing Office; 1981.
15. Bernat JL. How the distinction between "irreversible" and "permanent" illuminates circulatory-respiratory death determination. *J Med Philos*. 2010;35(3):242-255.
16. Repertinger S, Fitzgibbons WP, Omojola MF, Brumback RA. Long survival following bacterial meningitis-associated brain destruction. *J Child Neurol*. 2006;21(7):591-595.

Disputes & Debates: Rapid Online Correspondence

The editors encourage comments on recent articles through Disputes & Debates:

Access an article at [Neurology.org/N](https://www.neurology.org/N) and click on "MAKE COMMENT" beneath the article header.

Before submitting a comment to Disputes & Debates, remember the following:

- Disputes & Debates is restricted to comments about articles published in *Neurology* within 6 months of issue date, but the editors will consider a longer time period for submission if they consider the letter a significant addition to the literature
- Read previously posted comments; redundant comments will not be posted
- Your submission must be 200 words or less and have a maximum of 5 references; the first reference must be the article on which you are commenting
- You can include a maximum of 5 authors (including yourself)

Neurology®

Must Hypothalamic Neurosecretory Function Cease for Brain Death Determination?

No: The UDDA Revision Series

Panayiotis N. Varelas

Neurology 2023;101;137-139 Published Online before print July 10, 2023

DOI 10.1212/WNL.0000000000207336

This information is current as of July 10, 2023

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/101/3/137.full
References	This article cites 10 articles, 2 of which you can access for free at: http://n.neurology.org/content/101/3/137.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Brain death http://n.neurology.org/cgi/collection/brain_death Coma http://n.neurology.org/cgi/collection/coma Critical care http://n.neurology.org/cgi/collection/critical_care Endocrine http://n.neurology.org/cgi/collection/endocrine
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2023 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

