

# Neurology®

The most widely read and highly cited peer-reviewed neurology journal  
The Official Journal of the American Academy of Neurology



---

**Neurology Publish Ahead of Print**  
**DOI:10.1212/WNL.000000000206900**

Pearls & Oysters: Harnessing New Diagnostic and Therapeutic Approaches to Treat a Patient With Genetic Drug-Resistant Focal Epilepsy

**Author(s):**

Aafreen Khan, MBBS<sup>1</sup>; Erik H Middlebrooks, MD<sup>2</sup>; Pradeep Javarayee, MD<sup>3</sup>; William O Tatum, DO<sup>1</sup>; Sofia Sanchez Bolurate, MD, MSc<sup>1</sup>; Sanjeet S Grewal, MD<sup>4</sup>; Anteneh Feyissa<sup>1</sup>

**Corresponding Author:**

Anteneh Feyissa, feyissa.anteneh@mayo.edu

**Affiliation Information for All Authors:** 1. Department of Neurology, Mayo Clinic, Jacksonville, FL; 2. Department of Radiology, Mayo Clinic, Jacksonville, FL; 3. Department of Pediatric Neurology, Norton Children's Hospital, Louisville, KY; 4. Department of Neurosurgery, Mayo Clinic, Jacksonville, FL

**Equal Author Contribution:**

*Neurology*® Published Ahead of Print articles have been peer reviewed and accepted for publication. This manuscript will be published in its final form after copyediting, page composition, and review of proofs. Errors that could affect the content may be corrected during these processes.

**Contributions:**

Aafreen Khan: Drafting/revision of the manuscript for content, including medical writing for content; Major role in the acquisition of data; Analysis or interpretation of data

Erik H Middlebrooks: Drafting/revision of the manuscript for content, including medical writing for content; Major role in the acquisition of data; Analysis or interpretation of data

Pradeep Javarayee: Drafting/revision of the manuscript for content, including medical writing for content; Major role in the acquisition of data

William O Tatum: Drafting/revision of the manuscript for content, including medical writing for content

Sofia Sanchez Bolurate: Drafting/revision of the manuscript for content, including medical writing for content

Sanjeet S Grewal: Drafting/revision of the manuscript for content, including medical writing for content

Anteneh Feyissa: Drafting/revision of the manuscript for content, including medical writing for content; Major role in the acquisition of data; Study concept or design; Analysis or interpretation of data

**Figure Count:**

2

**Table Count:**

0

**Search Terms:**

[ 62 ] EEG, [ 66 ] Epilepsy surgery, [ 70 ] Epileptogenic zone, [ 75 ] Volumetric MRI use in epilepsy, [ 83 ] Cortical dysplasia

**Acknowledgment:****Study Funding:**

The authors report no targeted funding

**Disclosures:**

The authors report no relevant disclosures.

**Preprint DOI:****Received Date:**

2022-09-01

**Accepted Date:**

2022-12-22

**Handling Editor Statement:**

Submitted and externally peer reviewed. The handling editor was Resident and Fellow Section Editor Whitley Aamodt, MD, MPH.

**Abstract**

Focal cortical dysplasia (FCD) is a congenital developmental malformation and is one of the leading causes of drug-resistant focal epilepsy (DRFE). Although focal epilepsies traditionally have been regarded as acquired disorders, increasing evidence suggests a substantial genetic contribution to the pathogenesis of focal structural epilepsies, including FCDs. Mutations in the Dishevelled, Egl-10 and domain-containing protein 5 (DEPDC5) have recently emerged as a causative gene mutation in familial focal epilepsies associated with FCD type 2a, including Bottom of sulcus dysplasia (BOSD). We present a case of a 20-year-old man with DRFE, positive for DEPDC5 c.1555C>T (p.Gln519\*) heterozygous pathogenic variant. Initial 3T brain MRI was unrevealing, but subsequent 7T MRI including 7T edge-enhancing gradient echo (EDGE), revealed a left superior frontal sulcus BOSD concordant with the electroclinical data. The patient underwent treatment with MR-guided laser interstitial thermal ablation of the left frontal BOSD without intracranial EEG monitoring ("skipped candidate"), resulting in a seizure-free outcome of nine months since the last follow-up. Our case highlights the real-world application of summative information obtained through advancements in epilepsy–genetic testing, minimally invasive surgeries, and ultra-high field MRI, allowing us to provide a safe and effective treatment for a patient with a genetic DRFE.

## ***Pearls***

- Ultra-high field 7T MRI can enhance the detection of subtle anatomical lesions for MRI-negative epilepsy, including Bottom of sulcus dysplasia (BOSD), compared to 3T
- Increasing evidence suggests a substantial genetic contribution in the pathogenesis of focal structural epilepsies, including focal cortical dysplasias (FCDs)
- MR-guided laser interstitial thermal ablation therapy can safely and effectively treat drug-resistant focal epilepsy (DRFE) associated with a BOSD

## ***Oysters***

- Although genetic testing is infrequently performed in adults with lesional focal epilepsies, routine diagnostic testing is highly relevant in adults with childhood-onset nonlesional DRFE, or strong family history of epilepsy
- Identification of BOSD using advanced imaging techniques, including 7T MRI, could potentially prevent unnecessary invasive intracranial EEG evaluation

## Case Report

A 20-year-old left-handed student presented with drug-resistant epilepsy (DRE). He has a family history of epilepsy, with two maternal cousins having a seizure disorder. One cousin reportedly had absence seizures, while the other had tonic-clonic seizures. His seizures started at seven years of age. Initially, he had nocturnal bilateral tonic-clonic seizures, with subsequent development of daytime seizures characterized by an aura of a "fuzzy feeling in his brain" and right-sided "leg sensitivity." These would typically evolve into a focal aware motor seizure with vocalization ("no, no," "whoa"), grunting, and right-sided facial twitching. At times, these evolved into a right-sided head deviation and stiffening accompanied by loss of awareness. Infrequently, these would culminate in a focal to a bilateral tonic-clonic seizure. While the focal motor seizures occurred daily, the focal to bilateral tonic-clinic seizures occurred every other month. Seizures were refractory to treatment with four antiseizure medications (ASMs): oxcarbazepine, lamotrigine, brivaracetam, and clobazam. He was also failed by treatment with levetiracetam and valproic acid.

At age 18 years, he underwent presurgical evaluation at an outside hospital. Inpatient video EEG monitoring revealed rare low-to-moderate amplitude spike-and-wave discharges in the bifrontal head region (left > right). Multiple electroclinical seizures were also captured. EEG onset was poorly localizing, although most were predominantly associated with rhythmic alpha discharges in the left frontal lobe. Clinically, these were accompanied by right-sided facial twitch and repetitive mouth movements, perhaps repeating words. 3T brain MRI and PET was unrevealing.

Magnetoencephalography identified dipoles in the left superior frontal gyrus (**Figure 1, A–B**). Neuropsychological testing showed overall superior intelligence. Given his age of onset and strong family history of epilepsy, genetic testing was performed using an epilepsy gene panel (Invite, San Francisco, CA). This identified DEPDC5 c.1555C>T (p.Gln519\*) heterozygous pathogenic variant. His case was discussed at an epilepsy surgical conference and recommendation was made to proceed with intracranial EEG monitoring using a stereoelectroencephalography approach.

He underwent a repeat presurgical evaluation at our institution. Inpatient video-EEG revealed repetitive sharp, spikes, and polyspike waves in the left frontocentral and midline head regions (**Figure 1C**). Ictal EEG showed sustained bursts of fast beta activity, sharp waves, and poly spikes, most prominent in the vertex and left frontocentral head regions (**Figure 1D**). During these, he reported feelings of "right leg sensitivity" and facial twitching. He was aware and repeatedly vocalized "oh" and "no." An ultra-high field 7T brain MRI revealed a bottom of sulcus dysplasia (BOSD) in the left superior frontal sulcus, most evident on the 3D Edge-Enhancing Gradient Echo (EDGE) sequence (**Figure 2**). He was presented at our multidisciplinary epilepsy surgical conference. Recommendation was made to perform MRI-guided laser interstitial thermal ablation therapy (MRLiTT) on the left superior frontal sulcus BOSD. Given the concordant electro-clinical-radiological findings, intracranial EEG monitoring was deemed unnecessary. Following the MRLiTT, he has become seizure free for 9-months with stable doses of brivaracetam, clobazam, lamotrigine, and oxcarbazepine.

## DISCUSSION

We present a case of a young man with DRFE on whom a genetic test was performed, given his age of onset and strong family history of epilepsy. Identifying a DEPDC5 heterozygous pathologic variant prompted the pursuit of ultra-high field 7T-MRI that revealed a left superior frontal sulcus BOSD. The concordant electro-clinical-radiological findings obviated the need for invasive intracranial EEG monitoring. MRLiTT of the presumed BOSD leads to seizure-freedom, albeit he is yet to attain the ideal 2-year seizure-freedom cutoff.

Focal cortical dysplasias (FCDs) are localized regions of the malformed cerebral cortex and are frequently associated with epilepsy. Seizures associated with FCD are often resistant to ASMs.<sup>1</sup> In 2011, the ILAE Commissioned Task Force proposed a three-tiered classification system for FCDs. Briefly, FCD Type I refers to isolated lesions with dyslamination of the neocortex, Type II consists of isolated lesions characterized by cortical dyslamination and dysmorphic neurons, either without balloon cells (Type IIa) or with balloon cells (Type IIb), and Type III lesions are defined by their association with another principal lesion (e.g., mesial temporal lobe sclerosis).<sup>2</sup> BOSD is a highly localized form of FCD Type II in which the electrophysiologic, MRI, and pathologic abnormalities are confined to a single sulcus, maximal at the Bottom of the sulcus.<sup>3</sup>

Cortical thickening, subcortical hyperintensity, blurring in the gray-white matter interface, and transmantle sign are considered major findings in the MRI diagnosis of FCD. However, many of these features are more common in the more conspicuous Type IIb FCD.<sup>4</sup> Other signs include elongation and straightening of the dysplastic sulcus and depression or unusual sulcation of the overlying cerebral surface.<sup>4</sup> Recently, Tang et al. reported a "black line" sign on the 7T weighted gradient echo sequence in 20 patients with DRFE with pathologically confirmed FCD, but exclusively seen in Type IIb FCD.<sup>5</sup> FCD Type I and IIa are often far more subtle and frequently undetectable with conventional MRI. Besides, BOSDs, which can be Type IIa or IIb, typically found in the frontal and parietal lobes, are most likely to be overlooked.<sup>6</sup>

Advanced 7T MRI sequences have been shown to offer a superior contrast-to-noise ratio in detecting previously unseen subtle lesions of FCDs. In a recent study, 67 patients with DRFE and nonlesional 3T MRI reported that unaided visual review of 7T detected previously unappreciated subtle lesions in 22%; when aided by MRI post-processing, the yield increased to 43%.<sup>7</sup> However, morphometric analysis techniques have shown peak performance in detecting Type IIb FCD with poor detection of the more subtle Type I FCD. We show the value of 3D-EDGE at 7T in detecting subtle FCD. In the majority, the lesion location identified by 7T was identical or contained within the invasive EEG ictal onset zone. Complete resection of the 7T identified lesion resulted in a seizure-free outcome, with histopathology consisted mainly of FCD. In another study, 7T MRI detected new lesions in over a third of 3T MRI nonlesional patients, confirmed and better characterized a 3T suspected lesion in one-third of patients, and helped



exclude a 3T suspected lesion in the remainder.<sup>8</sup> Our case illustrates that the inherently low signal-to-noise ratio 3D-EDGE sequence greatly benefits from the added signal obtained at ultra-high field MRI, such as 7T, and may be one of the more valuable sequences in detecting both Type I and Type II FCD.<sup>9</sup> In centers where ultra-high field 7T MRI is unavailable, 3D-EDGE and other techniques, including a three-dimensional fluid-attenuated inversion recovery (3D FLAIR) and proton density MRI can be utilized to improve the detection of subtle FCDs.<sup>10</sup>

Increasing evidence suggests a strong genetic contribution to the pathogenesis of focal structural epilepsies, including FCDs.<sup>11</sup> For example, mutations in the Dishevelled, Egl-10 and domain-containing protein 5 (DEPDC5), a GTPase-activating protein, have recently emerged as a significant gene mutated in familial focal epilepsies associated with FCD IIa. Although genetic testing among children with epilepsy has become a part of routine testing, few studies have focused on genetic testing in adults.<sup>12</sup> A European study found a diagnostic yield of 23% with genetic testing in a cohort of adult patients with DRE.<sup>4</sup> In contrast, a more recent study from the US only found 10.9% of diagnostics with genetic testing.<sup>13</sup> Nevertheless, genetic testing of adults with focal epilepsies can help guide clinical management decisions. The diagnostic finding could have clinical implications beyond epilepsy care, including counseling for family planning. That said, the cost of testing, delays in receiving results, and problems interpreting results are barriers to its routine use in adults with focal epilepsy. Therefore, selecting the appropriate patient population that requires diagnostic genetic testing is crucial. The

yield is higher among patients with epileptic encephalopathies, childhood-onset and a strong family history of epilepsy, and nonlesional cases.

Identifying structural lesions on high-resolution MRI can obviate the need for invasive intracranial EEG monitoring, as in our patient. Ideal “skip candidates” for a one-stage surgery typically have concordant clinical data, including semiology, EEG, and a characteristic epileptogenic lesion on MRI. Traditionally open resection surgery has been the principal therapeutic option for treating FCD-associated DRFE. Newer minimally invasive therapies such as MRLiTT have also been reported in patients with FCDs.<sup>14</sup> Similar to our case, previous studies had undergone one-stage MRLiTT when the BOSD was distant from the eloquent cortex.<sup>15</sup>

This case sums up the cumulative impact of advancements in epilepsy–genetics, ultra-high field 7T MRI, and minimally invasive surgeries in allowing us to provide a safe and effective treatment for a patient with a genetic DRFE.

## Appendix 1. Authors

<b>Name</b>	<b>Location</b>	<b>Contribution</b>
Aafreen Khan, M.B.B.S.	Department of Neurology, Mayo Clinic, Jacksonville, FL	Drafting and revision of the manuscript for intellectual content, acquisition of data
Erik H. Middlebrooks, MD	Department of Radiology, Mayo Clinic, Jacksonville, FL	Drafting and revision of the manuscript for intellectual content, acquisition of data
Pradeep Javarayee, MD	Department of Pediatric Neurology, Norton Children's Hospital, Louisville, KY	Revised the manuscript for intellectual content and acquisition of data
William O. Tatum, DO	Department of Neurology, Mayo Clinic, Jacksonville, FL	Revised the manuscript for intellectual content
Sofia Sanchez Bolurate, MD, MSc	Department of Neurology, Mayo Clinic, Jacksonville, FL	Revised the manuscript for intellectual content
Sanjeet S. Grewal, MD	Department of Neurosurgery, Mayo Clinic, Jacksonville, FL	Revised the manuscript for intellectual content

Anteneh M. Feyissa, MD, MSc	Department of Neurology, Mayo Clinic, Jacksonville, FL	Design and conceptualized study; drafting and revision of the manuscript for intellectual content, acquisition of data
-----------------------------------	--	---

## References

1. Baldassari S, Ribierre T, Marsan E, et al. Dissecting the genetic basis of focal cortical dysplasia: a large cohort study. *Acta Neuropathol (Berl)*. 2019;138(6):885-900. doi:10.1007/s00401-019-02061-5
2. Blümcke I, Thom M, Aronica E, et al. The clinicopathologic spectrum of focal cortical dysplasias: A consensus classification proposed by an ad hoc Task Force of the ILAE Diagnostic Methods Commission<sup>1</sup>: The ILAE Classification System of FCD. *Epilepsia*. 2011;52(1):158-174. doi:10.1111/j.1528-1167.2010.02777.x
3. Harvey AS, Mandelstam SA, Maixner WJ, et al. The surgically remediable syndrome of epilepsy associated with bottom-of-sulcus dysplasia. *Neurology*. 2015;84(20):2021-2028. doi:10.1212/WNL.0000000000001591
4. Johannesen KM, Nikanorova N, Marjanovic D, et al. Utility of genetic testing for therapeutic decision-making in adults with epilepsy. *Epilepsia*. 2020;61(6):1234-1239. doi:10.1111/epi.16533


5. Tang Y, Blümcke I, Su TY, et al. Black Line Sign in Focal Cortical Dysplasia IIB: A 7T MRI and Electroclinicopathologic Study. *Neurology*. 2022;99(6):e616-e626. doi:10.1212/WNL.0000000000200702
6. Liu Z, Hu W, Sun Z, et al. MRI Abnormalities Predominate in the Bottom Part of the Sulcus with Type II Focal Cortical Dysplasia: A Quantitative Study. *Am J Neuroradiol*. 2019;40(1):184-190. doi:10.3174/ajnr.A5919
7. Wang I, Oh S, Blümcke I, et al. Value of 7T MRI and post-processing in patients with nonlesional 3T MRI undergoing epilepsy presurgical evaluation. *Epilepsia*. 2020;61(11):2509-2520. doi:10.1111/epi.16682
8. Bubrick EJ, Gholipour T, Hibert M, Cosgrove GR, Stufflebeam SM, Young GS. 7T versus 3T MRI in the presurgical evaluation of patients with drug-resistant epilepsy. *J Neuroimaging*. 2022;32(2):292-299. doi:10.1111/jon.12958
9. Middlebrooks EH, Lin C, Westerhold E, et al. Improved detection of focal cortical dysplasia using a novel 3D imaging sequence: Edge-Enhancing Gradient Echo (3D-EDGE) MRI. *NeuroImage Clin*. 2020;28:102449. doi:10.1016/j.nicl.2020.102449
10. Saini J, Singh A, Kesavadas C, et al. Role of three-dimensional fluid-attenuated inversion recovery (3D FLAIR) and proton density magnetic resonance imaging for the detection and evaluation of lesion extent of focal cortical dysplasia in patients with refractory epilepsy. *Acta Radiol*. 2010;51(2):218-225. doi:10.3109/02841850903433805

11. Baulac S, Ishida S, Marsan E, et al. Familial focal epilepsy with focal cortical dysplasia due to *DEPDC5* mutations: FCD and *DEPDC5* Mutations. *Ann Neurol*. 2015;77(4):675-683. doi:10.1002/ana.24368
12. Lee WS, Stephenson SEM, Pope K, et al. Genetic characterization identifies bottom-of-sulcus dysplasia as an mTORopathy. *Neurology*. 2020;95(18):e2542-e2551. doi:10.1212/WNL.0000000000010670
13. McKnight D, Bristow SL, Truty RM, et al. Multigene Panel Testing in a Large Cohort of Adults With Epilepsy: Diagnostic Yield and Clinically Actionable Genetic Findings. *Neurol Genet*. 2022;8(1):e650. doi:10.1212/NXG.0000000000000650
14. Guerrini R, Barba C. Focal cortical dysplasia: an update on diagnosis and treatment. *Expert Rev Neurother*. 2021;21(11):1213-1224. doi:10.1080/14737175.2021.1915135
15. Jain P, Ochi A, McInnis C, et al. Surgical outcomes in children with bottom-of-sulcus dysplasia and drug-resistant epilepsy: a retrospective cohort study. *J Neurosurg Pediatr*. 2021;28(3):295-305. doi:10.3171/2021.2.PEDS20967

## Figure Legends


**Figure 1 Title:** MEG images and EEG tracings show a left frontal seizure focus.

**Figure 1.** MEG source localization with the equivalent current model, sagittal (A) and coronal views (B), shows dipoles in the left superior frontal gyrus and insula (not shown). Interictal (C) and ictal (D) EEG tracings in a referential montage show left frontocentral and midline spikes and fast activity (arrowheads) and low amplitude fast seizure discharge arising from the same region (arrows-onset), respectively.



**Figure 2 Title:** Ultra-high field 7T brain MRI shows a bottom of sulcus dysplasia in the left superior frontal sulcus.

**Figure 2.** Sagittal 3D Edge-Enhancing Gradient Echo (3D-EDGE) acquired at 7T shows thickening and blurring of the normal junctional line at the gray matter-white matter interface (arrow) consistent with focal cortical dysplasia (A). Coronal 7T MP2RAGE T1-weighted uniform (UNI) image shows the blurring of the gray-white boundary (arrow) [B]. Postoperative coronal T1-weighted post-contrast image showing the enhancing ablation zone covering the area of FCD (C).





# Neurology®

## **Pearls & Oy-sters: Harnessing New Diagnostic and Therapeutic Approaches to Treat a Patient With Genetic Drug-Resistant Focal Epilepsy**

Aafreen Khan, Erik H Middlebrooks, Pradeep Javarayee, et al.

*Neurology* published online January 25, 2023

DOI 10.1212/WNL.0000000000206900

**This information is current as of January 25, 2023**

<b>Updated Information &amp; Services</b>	including high resolution figures, can be found at: <a href="http://n.neurology.org/content/early/2023/01/25/WNL.0000000000206900.full">http://n.neurology.org/content/early/2023/01/25/WNL.0000000000206900.full</a>
<b>Subspecialty Collections</b>	This article, along with others on similar topics, appears in the following collection(s): <b>Cortical dysplasia</b> <a href="http://n.neurology.org/cgi/collection/cortical_dysplasia">http://n.neurology.org/cgi/collection/cortical_dysplasia</a> <b>EEG</b> <a href="http://n.neurology.org/cgi/collection/eeg_">http://n.neurology.org/cgi/collection/eeg_</a> <b>Epilepsy surgery</b> <a href="http://n.neurology.org/cgi/collection/epilepsy_surgery_">http://n.neurology.org/cgi/collection/epilepsy_surgery_</a> <b>Epileptogenic zone</b> <a href="http://n.neurology.org/cgi/collection/epileptogenic_zone">http://n.neurology.org/cgi/collection/epileptogenic_zone</a> <b>Volumetric MRI use in epilepsy</b> <a href="http://n.neurology.org/cgi/collection/volumetric_mri_use_in_epilepsy">http://n.neurology.org/cgi/collection/volumetric_mri_use_in_epilepsy</a>
<b>Permissions &amp; Licensing</b>	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: <a href="http://www.neurology.org/about/about_the_journal#permissions">http://www.neurology.org/about/about_the_journal#permissions</a>
<b>Reprints</b>	Information about ordering reprints can be found online: <a href="http://n.neurology.org/subscribers/advertise">http://n.neurology.org/subscribers/advertise</a>

*Neurology*® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2023 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

